



Mycoplasma Bovis Bacterin



Cattle Vaccine

100 Dose / 200 mL For Animal Use Only

Mycoplasma Bovis Bacterin

Indications: This product has been shown to be effective for the vaccination of healthy cattle four months of age or older against Mycoplasma Bovis. Duration of immunity has not been determined. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Directions: Mix well and aseptically inject 2.0 mL subcutaneously in the side of the neck. Repeat on days 7 and 14 alternating sides of the neck.

Cautions: Do not mix with other products. May cause local swelling. Do not inject into muscle, may cause carcass trim. Do not vaccinate within 60 days of slaughter. Do not contaminate with dirt or chemicals. Use sterile equipment and adequately clean and disinfect bottle cap prior to each entry. Use entire contents when opened. Store at 2° to 8° C. Do not use if emulsion is broken as indicated by separation at the BOTTOM of bottle. In case of human exposure, contact a physician. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy. This product has not been tested in pregnant animals.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.

***Adjuvant** – Bimeda Biological's Proprietary Technology.



8 99855 10045 0

VLN / PCN: 290 / 2760.01



100 Dose / 200 mL

Bimeda Biologicals, Inc.
San Angelo, Texas
(800) 284-8403

Exp. Date:

Serial Number: